

Effect of co-extraction of pomegranate seed oil with green tea leaves on the extraction yield and quality of extracted oil

Zeinab Dezashibi¹, Sodeif Azadmard-Damirchi^{1,*}  and Zahra Piravi-Vanak²

¹ Department of Food Science and Technology, Faculty of Agriculture, University of Tabriz, Tabriz, Iran

² Food, Halal, and Agricultural Products Research Group, Food Technology and Agricultural Products Research Center, Standard Research Institute (SRI), Karaj, Iran

Received 28 January 2022 – Accepted 13 May 2022

Abstract – Co-extraction of pomegranate seed oil (PSO) with green tea leaves (GTL) (0 [control sample], 2.5, 5, 7.5, and 10% w/w) was conducted by cold-press to evaluate the extracted oil quality during storage. The extraction yield was similar to the control sample up to 5% w/w of GTL. Total phenol and chlorophyll content were increased in the extracted oils with increasing the level of GTL. The acid and peroxide values were increased during the storage and the lowest values were achieved for PSO with 5% GTL. Rancimat analysis also confirmed the PV results, therefore using 5% GTL led to providing the highest induction period (11.5 h). Consequently, using 5% of GTL during the extraction of PSO by cold-press could result in an appropriate extraction yield and also present higher oxidation stability. This method does not need any antioxidant extraction from herbs and is very cost-effective, time-efficient, and uses no chemicals.

Keywords: oil extraction / oxidative stability / quality / cold-press

Résumé – Effet de la co-extraction de l'huile de pépins de grenade avec des feuilles de thé vert sur le rendement d'extraction et la qualité de l'huile extraite. La co-extraction de l'huile de pépins de grenade (PSO) avec des feuilles de thé vert (GTL) (0 – échantillon témoin – , 2,5, 5, 7,5, et 10 % p/p) a été réalisée par pression à froid pour évaluer la qualité de l'huile extraite durant son stockage. Le rendement d'extraction était similaire à celui de l'échantillon témoin jusqu'à 5 % p/p de GTL. La teneur en polyphénols totaux et en chlorophylle a augmenté dans les huiles extraites avec l'augmentation du taux de GTL. Les indices d'acide et de peroxyde (PV) ont augmenté pendant le stockage et les valeurs les plus basses ont été obtenues pour la PSO avec 5 % de GTL. L'analyse Rancimat a également confirmé les résultats du PV. Ainsi, l'utilisation de 5 % de GTL a conduit à fournir la période d'induction la plus longue (11,5 h). Par conséquent, l'utilisation de 5 % de GTL pendant l'extraction de PSO par pression à froid pourrait offrir un rendement d'extraction optimal et présenter également une plus grande stabilité à l'oxydation. Cette méthode ne nécessite pas d'extraction d'antioxydants à partir de plantes et est très rentable, efficace en termes de temps et n'utilise aucun produit chimique.

Mots clés : extraction d'huile / stabilité à l'oxydation / qualité / pression à froid

Highlights

- Pomegranate seed oil has many applications in the pharmaceutical, cosmetic and food industries.
- Co-extraction of pomegranate seed oil with green tea leaves was conducted by cold-press.
- The oil extraction yield was not affected up to 5% w/w of green tea leaves incorporation.

- Incorporating green tea leaves during pomegranate seed oil extraction could produce high-quality oil.
- This method is very cost-effective, time-efficient and uses no chemicals.

1 Introduction

Pomegranate (*Punica granatum* L.) has been consumed since ancient times due to its nutritional value (Khoddami *et al.*, 2014). It has been originated in the Himalayas and Mediterranean region and has mostly grown in tropical and

*Corresponding author: sodeifazadmard@yahoo.com

subtropical countries such as India, Iran, China, and the USA (Khoddami *et al.*, 2014; Paul and Radhakrishnan, 2020).

Pomegranate seed is accounted as a by-product of the pomegranate juice industry. Pomegranate seeds are a rich source of oil corresponding to 12–20% of total seed weight (Costa *et al.*, 2019; Drinić *et al.*, 2020). Pomegranate seed oil (PSO) contains many valuable components including vitamin E, sterols, and more important the “punicic acid”. PSO has a remarkable effect on health due to its high content punicic acid (18:3, 9-*cis*, 11-*trans*, 13-*cis*) varies from 32–87%. Other main fatty acid of PSO are C18:2 (up to 24%) followed by C18:1 (up to 17%), 18:0 (up to 2.6%) and 16:0 (up to 4.6%) (Khoddami *et al.*, 2014). Beneficial properties of PSO have changed the pomegranate juice residues from low-value by-products (mainly as waste) to high valuable biomaterial (Khoddami *et al.*, 2014; Tian *et al.*, 2013). Today, PSO has medicinal, health, and food uses especially for its punicic acid, which has pharmaceutical effects (de Melo *et al.*, 2016). However, PSO, due to its high content of unsaturated fatty acid profiles, is sensitive to oxidation (Costa *et al.*, 2019; Drinić *et al.*, 2020; Rezvankhah *et al.*, 2022).

Oil oxidative stability can be improved using antioxidants being added after extraction (Drinić *et al.*, 2020). Antioxidants have several types and mainly they are divided into two synthetic and natural ones. There are concerns about the consumption of synthetic antioxidants due to their potential adverse health effects (Drinić *et al.*, 2020).

Recently, attention to the use of natural antioxidants has increased, and especially consumers prefer to use natural antioxidants instead of synthetic ones (Drinić *et al.*, 2020). The oils can be enriched with natural antioxidants and, thus, the oxidation and rancidity are retarded and oil quality is improved during storage (Osanloo *et al.*, 2021).

Plants are rich in polyphenolic compounds, which potentially indicate antioxidant activity and radical scavenging activity. Green tea leaves (GTL) are one of the richest sources of polyphenolic compounds (Li and Jiang, 2010; Musial *et al.*, 2020).

There are many options, such as using antioxidants, extraction of oils from mixed seedoils, and pretreatments of seedoils before oil extraction, to enhance oxidative stability of oils. Dehghan-Manshadi *et al.* (2020) applied infrared (IR)-assisted spouted bed drying (SBD) as a potential alternative to the traditional hot air drying for heat-sensitive components of flaxseeds. Mazaheri *et al.* (2019a) reported that microwave treatment of black seed before the oil extraction reduced lipase activity more effectively. They reported that the treatment of black seeds with microwaves and soaking them before extracting oil in a cold-press raised the oil oxidative stability. Nain *et al.* (2021) reported that using green tea enhanced the oxidative stability of DHA-rich oil by reducing the formation of peroxides and secondary oxidation products. Drinić *et al.* (2020) used pomegranate peel extract in PSO individually or with a combination with BHT to enhance the oxidative stability of the oil. Mazaheri *et al.* (2019b) investigated the cold-press extraction of sunflower and black cumin seeds and reported that seeds mixing caused an increase in oxidative stability, which increased from 6.78 to 9.69 h.

All of the above-mentioned ways to enhance extracted oils' oxidative stability have advantages and disadvantages. One new way can be blending seedoils with plant materials high in bioactive compounds such as GTL and their

coextraction by the press. Therefore, the aim of this work is the enhancement of PSO oxidative stability using GTL as a natural antioxidant. Co-extraction of antioxidant compounds of GTL with PSO as a new method of extraction can be highlighted in industrial applications concerning the production of oil with improved oxidative stability and high content of bioactive components.

2 Materials and method

2.1 Materials

Pomegranates seeds, which were separated from the Iranian pomegranate cultivar (Saveh Sweet White Peel) after juice extraction, were purchased from local market (Tabriz, Iran). Green tea leaves (*Camellia sinensis* L.) were obtained from local tea market (Tabriz, Iran). All chemical materials were provided from Sigma-Aldrich Co. (St. Louis, Missouri, United States).

2.2 Extraction process

PSO was coextracted with green tea leaves using a mechanical screw press apparatus (model P500R, Anton Fries, Germany) (Piravi-Vanak *et al.*, 2022; Rostami *et al.*, 2014). First, pomegranate seeds and green tea leaves were conditioned to moisture contents of 6.4% and 3.7% based on the preliminary oil extraction yield. Then, the mixture of PSO and GTL at a level of 0 (control sample), 2.5, 5, 7.5, and 10% (w/w) simultaneously undergone the extraction process by screw press apparatus. The temperature of extracted oil was not raised over 40 °C during the oil extraction by cold-press. Extracted oil samples were kept for 90 days at room condition in a dark place and their quality was analyzed on the extraction day and every 30 days of storage.

Extraction yield

The extraction yield (EY) was calculated based on the weight of obtained oil after extraction and the weight of pomegranate seed powder as the following equation (Rezvankhah *et al.*, 2018):

$$\text{Extraction yield (\%)} = \frac{\text{Extracted oil amount (g)}}{\text{initial seed powder (g)}} \times 100 \quad (1)$$

2.3 Determination of total phenolic content

The total phenolic content of PSO enriched with GTL was determined using the Folin-Ciocalteu method according to the method of Womeni *et al.* (2016) with brief modification. Firstly, 1 mL of oil samples (1 mg/mL) that dissolved in ethanol was mixed with 1 mL of diluted Ciocalteu reagent (1:10, v/v) and 3 mL of sodium carbonate 7%. The prepared mixtures were kept in a dark place at an ambient temperature for 30 min. The absorbance of oil mixtures was determined at 760 nm using a UV-Vis spectrophotometer. Also, pure ethanol was used as the control, and gallic acid at the concentration of 0–0.06 mg/mL was used as a standard. Eventually, the TPC of oil mixtures was expressed as milligram gallic acid equivalents (GAE)/gram samples.

2.4 Determination of chlorophyll

Chlorophyll content of PSO enriched with GTL was determined at 670 nm in cyclohexane using the method of [Sahin *et al.* \(2017\)](#). The chlorophyll concentration was expressed as mg of pheophytin per kg of oil.

2.5 Determination of acid value (AV)

AV was determined by potentiometric titration in an automatic titrator model G20 (Mettler Toledo, Urdorf, Switzerland) according to [Costa *et al.* \(2019\)](#). Briefly, oil acidity was analyzed by titrating PSO enriched with GTL dissolved in 45 mL acetone: ethanol (1:1, v/v) with 0.4 M NaOH until pH 11. AV was expressed as % punnic acid.

2.6 Determination of peroxide value (PV)

The PV was measured according to the method described by [Rezvankhah *et al.* \(2019\)](#). Determination was conducted by titration of 0.1 N KI saturated solutions of the oil with 0.1 N Na₂S₂O₃ and starch as an indicator. PV was expressed in mg oxygen equivalent per kg of oil (meqO₂/kg-oil).

2.7 Rancimat analysis

The oxidative stability of PSO before and after enrichment was evaluated by the Rancimat method at 110 °C in a Rancimat apparatus (Metrohm 743; Metrohm Co., Herisau, Switzerland) according to [Costa *et al.* \(2019\)](#). The oxidation process was conducted upon 2.5 g oil samples at the mentioned temperature at an air velocity of 20 L/h while monitoring mixtures until a sharp increase in the water conductivity corresponded to the oil stability index. The stability of oil mixtures was expressed as the induction period (IP).

2.8 Statistical analysis

All data were conducted in three replications and provided in mean and standard deviation. The statistical analysis was conducted by one-way ANOVA and the mean difference between the data was implemented by the Duncan test at the probable level of 5% ($P < 0.05$) using SPSS software (IBM SPSS Statistics 23).

3 Results and discussion

3.1 Extraction yield

The extraction yield in the cold-press extraction method is affected mainly by the temperature and also moisture content of the seed ([Naebi *et al.*, 2022](#)). The effects of the incorporation of green tea leaves on the extraction yield of PSO were evaluated during the mechanical press. Incorporation of GTL at a concentration ranging between 2.5 to 5% did not have a significant effect ($P < 0.05$) on the extraction yield (6.3 to 6.1%) ([Fig. 1](#)). However, incorporation of higher than 7.5% significantly reduced the extraction yield. It was thought that GTL at higher concentrations made a mechanical obstacle




Fig. 1. Extraction yield of pomegranate seed oil mixed with green tea leaves at the conditioned moisture content of 6.4% and 3.75%. T0, T2.5, T5, T7.5, and T10 indicate pomegranate seed (control sample), pomegranate seed with 2.5% green tea leaves, pomegranate seed with 5% green tea leaves, pomegranate seed with 7.5% green tea leaves, pomegranate seed with 10% green tea leaves, respectively.

against the elicitation of oil from the ground seed texture. [Eikani *et al.* \(2012\)](#) reported that the cold-pressing method indicated 4.29% extraction efficiency. Our extraction yield values were higher which can be attributed to the variety and also process condition.

It has been reported that a heating process by temperature 75 °C and moisture content of output seeds in the confine of 6.3–6.5% have yielded the highest extraction efficiency that resulted in high-quality oil production and by-products with the minimum remaining oil content according to the previous studies ([Rostami *et al.*, 2014](#)). Also, [Naebi *et al.* \(2022\)](#) declared that the optimum moisture content which gave the highest oil yield (33.5%) was 7.5%. The extracted PSO main fatty acids were punnic acid (C18:3) up to 76% as the main fatty acid followed by linoleic acid (C18:2) about 8.2%, oleic acid (C18:1) about 7.6%, palmitic acid (C16:0) about 4.6% and stearic acid (C18:0) about 2.6% ([Khoddami *et al.*, 2014](#)).

3.2 Total phenolic content

GTL is rich in antioxidants mainly catechins include epicatechin, epigallocatechin, epicatechin gallate, epigallocatechin gallate, and flavonoid compounds including caffeic acid, quercetin, hesperidin, and hesperetin that inherently increase the TPC ([Ebrahimi Monfared *et al.*, 2022](#); [Musial *et al.*, 2020](#); [Palit *et al.*, 2008](#); [Samadi and Raouf Fard, 2020](#); [Yeasmen and Orsat, 2021](#)). Although it has been reported that PSO has a high content of phenolic compounds ([Costa *et al.*, 2019](#); [de O Silva *et al.*, 2019](#)), the incorporation of GTL significantly enhanced the TPC of extracted PSO ([Tab. 1](#)). Phenolic acids, quercetin and naringenin have also been found in PSO ([Costa *et al.*, 2019](#)).

TPC ranged between 2.7 to 12.1 mg GAE/g for control and PSO with 10% incorporated GTL, respectively, which shows

Table 1. Total phenol content of extracted pomegranate seed oil incorporated with green tea leaves during the storage.

Sample	Storage time (day)			
	1	30	60	90
T0*	2.7 ^{Be**}	2.3 ^{Ce}	3.4 ^{Ad}	3.7 ^{Ad}
T2.5	3.4 ^{Bd}	2.8 ^{Cd}	5.9 ^{Ac}	6.1 ^{Ac}
T5	5.6 ^{Cc}	4.4 ^{Dc}	6 ^{Bc}	7.8 ^{Ab}
T7.5	8.2 ^{Ab}	4.7 ^{Cb}	5.3 ^{Bb}	8 ^{Ab}
T10	12.1 ^{Aa}	10 ^{Ba}	12.2 ^{Aa}	11.9 ^{Aa}

* For samples see [Figure 1](#).

** Different small superscripts indicate the significant difference between the samples while the different capital superscripts indicate the significant difference between the days ($P < 0.05$).

Table 2. Chlorophyll content of extracted pomegranate seed oil incorporated with green tea leaves during the storage.

Sample	Storage time (day)			
	1	30	60	90
T0*	0.18 ^{Ae**}	0.18 ^{Ad}	0.15 ^{ABe}	0.12 ^{Bc}
T2.5	2.1 ^{Ad}	2 ^{Ac}	1.7 ^{Bd}	1 ^{Cd}
T5	4.3 ^{Ac}	3.9 ^{Bb}	2.8 ^{Cc}	2.8 ^{Cc}
T7.5	5 ^{Ab}	4 ^{Bb}	3.3 ^{Cb}	3.5 ^{Cb}
T10	8.6 ^{Aa}	6.5 ^{Ba}	6 ^{Ca}	6 ^{Ca}

* For samples see [Fig. 1](#).

** Different small superscripts indicate the significant difference between the samples while the different capital superscripts indicate the significant difference between the days ($P < 0.05$).

the positive effect of GTL incorporation on the TPC on PSO. High TPC (10.44 mg GAE/g sample) content was reported for the PSO extracted by cold-pressing from the variety Torshe Malas Iran ([Khoddami et al., 2014](#)).

It has been reported that GTL has a high content of polyphenolic compounds, which can donate antioxidant activity ([Hu et al., 2021](#); [Musial et al., 2020](#); [Palit et al., 2008](#)). Catechins are the prominent polyphenols of GTL that can render antioxidant and preservation effects ([Palit et al., 2008](#); [Unno et al., 2018](#)). The addition of tea polyphenols to oil has shown other beneficial impacts ([Palit et al., 2008](#); [Samadi and Raouf Fard, 2020](#)). For instance, it has been reported that rice bran oil cold performance quality was improved after the addition of tea polyphenols. Indeed, rice bran oil retained clear at 0 °C for 5.5 h when tea polyphenols concentration was 0.08% w/w, which could recrystallize in rice bran oil before rice bran crystallization and further promote the nucleation of rice bran oil ([Wang et al., 2021](#)).

3.3 Chlorophyll content

The plants are rich in natural pigments such as carotenoids and chlorophylls. GTL is also rich in chlorophylls and incorporation of them enhances the chlorophyll content. The addition of GTL significantly ($P < 0.05$) increased the chlorophyll content of PSO on all days of storage ([Tab. 2](#)). It has been reported that extracted PSO by the supercritical CO₂ method did not have detectable chlorophyll ([de O Silva et al., 2019](#)). Storing had also a significant effect and chlorophyll

content was decreased during 90 days of storage. This reduction was associated with chlorophyll degradation ([Li et al., 2019](#); [Sahin et al., 2017](#)). It should be also pointed out that the addition of a high amount of chlorophyll to the oils with higher PUFAa has sensitizing effects and can increase oxidation ([Machado et al., 2022](#); [Rezvankhah et al., 2018, 2019](#); [Roshanak et al., 2016](#)). Chlorophylls can promote the initial autooxidation by their ability to generate alkyl hydroperoxides ([Machado et al., 2022](#); [Rezvankhah et al., 2018](#)). Based on current reports, the presence of high content of chlorophylls in the extracted oil can reduce the oil oxidative stability ([Rezvankhah et al., 2018, 2019](#)). So, the incorporation of a high amount of GTL into PSO can have adverse effects on oxidative stability.

3.4 Acid value

AV is associated with the free fatty acids and for PSO, punicic acid is principal ([Goula and Adamopoulos, 2012](#); [Keskin Cavdar et al., 2017](#); [Tian et al., 2013](#)). The higher the AV determination, the lower the quality of the obtained oil ([Drinić et al., 2020](#)). Degradation of triacylglycerol molecules especially occurs during storage due to the presence of oxidation parameters such as inducing moisture, metals, oxygen, etc. ([Rezvankhah et al., 2018, 2019, 2020](#)).

It was observed that incorporation of GTL especially at higher concentrations increased the AV ([Tab. 3](#)). Also, the incorporation of the GTL enhanced the AV during the storage time. The lowest amount of AV (0.1%) was obtained for the

Table 3. The acid value of extracted pomegranate seed oil incorporated with green tea leaves during the storage.

Sample	Storage time (day)			
	1	30	60	90
T0*	0.9Dc**	1.1 ^{Cd}	1.8 ^{Bc}	2.1 ^{Ab}
T2.5	0.9 ^{Dc}	1.2 ^{Cc}	1.5 ^{Bd}	1.8 ^{Ac}
T5	0.1 ^{Dc}	1.2 ^{Cc}	1.7 ^{Bc}	1.9 ^{Ac}
T7.5	1.3 ^{Db}	1.8 ^{Cb}	2.01 ^{Bb}	2.2 ^{Ab}
T10	1.5 ^{Ca}	2.1 ^{Ba}	2.3 ^{Aa}	2.4 ^{Aa}

* For samples see [Figure 1](#).

** Different small superscripts indicate the significant difference between the samples while the different capital superscripts indicate the significant difference between the days ($P < 0.05$).

Table 4. Peroxide value of extracted pomegranate seed oil incorporated with green tea leaves during the storage.

Sample	Storage time (day)			
	1	30	60	90
T0*	2.2Da**	3.7 ^{Ca}	4.9 ^{Ba}	5.2 ^{Aa}
T2.5	2.1 ^{Da}	2.3 ^{Cc}	2.7 ^{Bd}	2.9 ^{Ad}
T5	2.2 ^{Ba}	2.3 ^{Bc}	2.9 ^{Ac}	3 ^{Ad}
T7.5	2.3 ^{Ca}	2.4 ^{Cc}	3 ^{Bc}	3.4 ^{Ac}
T10	2.3 ^{Da}	2.8 ^{Cb}	3.5 ^{Bb}	4.7 ^{Ab}

* For samples see [Figure 1](#).

sample with 5% GTL on day 1 and the highest amount (2.4%) was obtained for PSO with 10% GTL on day 90. Drinić *et al.* (2020) reported that the AV was 0.84 mL/g, Dadashi *et al.* (2013) reported 3.78 and 8.36 mL/g for *n-hexane* extraction while 0.63 mL/g for cold-press technology (de Melo *et al.*, 2016).

3.5 Peroxide value

Although the incorporation of GTL increased the oil AV, the presence of phenolic compounds can reduce the oxidation rate associated with their antioxidant activity (Maqsood *et al.*, 2014). Also, PSO inherently has antioxidant activity mainly due to the presence of tocopherols and phytosterols (Costa *et al.*, 2019; de Melo *et al.*, 2016). According to Table 4, PV was determined after the incorporation of various concentrations of GTL and during storage. PV was decreased after the addition of 2.5% to PSO while higher concentration had even inducing effects on the oxidation of PUFAs (Bahmaei *et al.*, 2005; Palit *et al.*, 2008). The highest PV (4.7 meq-O₂/kg-oil) was obtained for PSO enriched with 10% GTL and the lowest amount (2.1 meq-O₂/kg-oil) was obtained for PSO with 2.5% GTL. The addition of a sufficient not excessive amount of polyphenol-rich source to the oils with high susceptibility to oxidation can retard the oxidation and inhibit the propagation (Mazaheri *et al.*, 2019a, 2019b; Palit *et al.*, 2008). However, the higher the polyphenol content, and also the higher the chlorophyll content the higher peroxy radical production, and thus, oxidation was increased (Li *et al.*, 2019; Rezvankhah *et al.*, 2019). Li *et al.* (2019) declared that residues of

chlorophyll induced oxidative rancidity and deterioration through photooxidation in rapeseed oil. They reported that PV of rapeseed oil exposed to light increased significantly, especially for the oil with a high content of chlorophyll. It was also consistent with reports by Bahmaei *et al.* (2005) that pigments present in oil impart an undesirable color and also promote oxidation in the presence of light. Regarding the storage, it was observed that PV was increased in all samples. However, the sample with 2.5% GTL indicated lower PV than control and other samples with higher GTL concentrations (Maqsood *et al.*, 2014). Incorporation of the high amount of GTL into the PSO-rich PUFAs leads to an increase in sensitivity of oil to oxidation since chlorophylls act as light adsorbents and, therefore, oil oxidation be propagated (Rezvankhah *et al.*, 2018, 2019). In addition, based on the current studies, the addition of herbs event without extraction can raise the oil oxidative stability (Ebrahimi Monfared *et al.*, 2022; Nain *et al.*, 2021). As reported by Ebrahimi Monfared *et al.* (2022), the addition of matcha GTL increased the oil oxidative stability led to a higher shelf-life of muesli. Also, Kırmızıyaya *et al.* (2021) reported that black, green, and white tea infusions and powder improved the oxidative stability of minced beef throughout refrigerated storage.

3.6 Rancimat

Plants due to their phenolic compounds can enhance the oxidative stability of oils. In the present study, the incorporation of GTL increased the oxidative stability of PSO (Fig. 2). Concerning this, IP values were determined and the lowest




Fig. 2. The induction period of pomegranate seed oil with and without green tea leaves at various concentrations (2.5 to 10%). T0, T2.5, T5, T7.5, and T10 indicate pomegranate seed (control sample), pomegranate seed with 2.5% green tea leaves, pomegranate seed with 5% green tea leaves, pomegranate seed with 7.5% green tea leaves, pomegranate seed with 10% green tea leaves, respectively.

amount was obtained for control and the highest amount was obtained for PSO with 5% GTL. Based on the PV and AV results, a higher concentration of GTL induces oxidation due to the high content of chlorophyll and degradation of phenol functional hydroxyl groups, which propagate the oxidation (Li *et al.*, 2019; Womeni *et al.*, 2016; Ye *et al.*, 2020). Thus, a moderate amount of GTL can be more effective in retarding the oxidation. Womeni *et al.* (2016) used old Cameroonian GTL and BHT to evaluate the oxidative stability of RBD palm olein under forced storage conditions. Based on the Rancimat results, the oil with GTL had a higher induction time (24.8–28.9 h) compared with BHT (20.1–22.7 h). Due to the presence of gallic acid, epicatechin gallate, galocatechin, and epigallocatechin gallate as the main phenolic antioxidants in GTL, the oils with these natural antioxidants would have higher oxidation stability (Womeni *et al.*, 2016). Mazaheri *et al.* (2019b) reported that co-extraction of sunflower oil with black seed oil increased the oxidative stability related to the presence of high phenolic compounds in black seed oil.

4 Conclusion

PSO lacking high oxidation stability was enriched with GTL at various concentrations (2.5 to 10%). Using up to 5% GTL did not affect the oil extraction yield. TPC as health-promoting components was increased in the PSO after the inclusion of GTL. In conclusion and general view and based on the factors related to the oxidative stability (AV, PV, and Rancimat analysis), it can be suggested that that PSO coextracted with 5% GTL has good quality and high oxidative stability. Therefore, the coextraction of oil from a mixture of plant materials and seed oils can be suggested as a safe, cost-effective, and time-efficient process for the industry.

Acknowledgments. This study was funded by the University of Tabriz, Tabriz, Iran.

Conflicts of interest. The authors declare that they have no conflicts of interest in relation to this article.

References

- Bahmaei M, Sadat Sabbaghian E, Farzadkish E. 2005. Development of a method for chlorophyll removal from canola oil using mineral acids. *Journal of the American Oil Chemists' Society* 82(9): 679–684.
- Barbosa DS. 2007. Green tea polyphenolic compounds and human health. *Journal für Verbraucherschutz und Lebensmittelsicherheit* 2(4): 407–413. <https://doi.org/10.1007/S00003-007-0246-Z>.
- Bharti R, Singh B. 2020. Green tea (*Camellia assamica*) extract as an antioxidant additive to enhance the oxidation stability of biodiesel synthesized from waste cooking oil. *Fuel* 262: 116658. <https://doi.org/10.1016/j.fuel.2019.116658>.
- Costa AMM, Silva LO, Torres AG. 2019. Chemical composition of commercial cold-pressed pomegranate (*Punica granatum*) seed oil from Turkey and Israel, and the use of bioactive compounds for samples' origin preliminary discrimination. *Journal of Food Composition and Analysis* 75: 8–16. <https://doi.org/10.1016/j.jfca.2018.09.004>.
- Dadashi S, Mousazadeh M, Emam-Djomeh Z, Mousavi SM. 2013. Pomegranate (*Punica granatum* L.) seed: a comparative study on biochemical composition and oil physicochemical characteristics. *International Journal of Advanced Biological and Biomedical Research* 1(4): 351–363.
- de Melo ILP, de Carvalho EBT, de Oliveira e Silva AM, *et al.* 2016. Characterization of constituents, quality and stability of pomegranate seed oil (*Punica granatum* L.). *Food Science and Technology* 36(1): 132–139. <https://doi.org/10.1590/1678-457X.0069>.
- de O Silva L., Ranquine L.G., Monteiro M., Torres A.G. 2019. Pomegranate (*Punica granatum* L.) seed oil enriched with conjugated linolenic acid (cLnA), phenolic compounds and tocopherols: improved extraction of a specialty oil by supercritical CO₂. *The Journal of Supercritical Fluids* 147: 126–137. <https://doi.org/10.1016/J.SUPFLU.2019.02.019>.
- Dehghan-Manshadi A, Peighambaroust SH, Azadmard-Damirchi S, Niakousari M. 2020. Effect of infrared-assisted spouted bed drying of flaxseed on the quality characteristics of its oil extracted by different methods. *Journal of the Science of Food and Agriculture* 100(1): 74–80. <https://doi.org/10.1002/JSFA.9995>.
- Drinić Z, Mudrić J, Zdunić G, Bigović D, Menković N, Šavikin K. 2020. Effect of pomegranate peel extract on the oxidative stability of pomegranate seed oil. *Food Chemistry* 333: 127501. <https://doi.org/10.1016/j.foodchem.2020.127501>.
- Ebrahimi Monfared K, Gharachorloo M, Jafarpour A, Varvani J. 2022. Production feasibility of functional probiotic muesli containing matcha and investigation of its physicochemical, microbial, and sensory properties. *Journal of Food Measurement and Characterization* 16: 975–986.
- Eikani MH, Golmohammad F, Homami SS. 2012. Extraction of pomegranate (*Punica granatum* L.) seed oil using superheated hexane. *Food and Bioproducts Processing* 90(1): 32–36. <https://doi.org/10.1016/J.FBP.2011.01.002>.
- Goula AM, Adamopoulos KG. 2012. A method for pomegranate seed application in food industries: seed oil encapsulation. *Food and Bioproducts Processing* 90(4): 639–652. <https://doi.org/10.1016/J.FBP.2012.06.001>.

- Hu L, Li T, Luo Q, Zhang J. 2021. Antioxidant stability of colloidal tea polyphenols in tea seed oil. *Journal of Food Processing and Preservation* 45(2): e15130. <https://doi.org/10.1111/JFPP.15130>.
- Keshavarzi M, Najafi G, Ahmadi Gavlighi H, Seyfi P, Ghomi H. 2020. Enhancement of polyphenolic content extraction rate with maximal antioxidant activity from green tea leaves by cold plasma. *Journal of Food Science* 85(10): 3415–3422. <https://doi.org/10.1111/1750-3841.15448>.
- Keskin Çavdar H, Koçak Yanık D, Gök U, Göğüs F. 2017. Optimisation of microwave-assisted extraction of pomegranate (*Punica granatum* L.) seed oil and evaluation of its physicochemical and bioactive properties. *Food Technology and Biotechnology* 55(1): 86–94. <https://doi.org/10.17113/FTB.55.01.17.4638>.
- Khoddami A, Man YBC, Roberts TH. 2014. Physico-chemical properties and fatty acid profile of seed oils from pomegranate (*Punica granatum* L.) extracted by cold-pressing. *European Journal of Lipid Science and Technology* 116(5): 553–562. <https://doi.org/10.1002/EJLT.201300416>.
- Kırmızızkaya G, Karakaya M, Babaoğlu AS. 2021. Black, green, and white tea infusions and powder forms improve oxidative stability of minced beef throughout refrigerated storage. *Journal of Food Processing and Preservation* 45(4): e15359. <https://doi.org/10.1111/jfpp.15359>.
- Li DC, Jiang JG. 2010. Optimization of the microwave-assisted extraction conditions of tea polyphenols from green tea. *International Journal of Food Sciences and Nutrition* 61(8): 837–845. <https://doi.org/10.3109/09637486.2010.489508>.
- Li X, Yang R, Lv C, et al. 2019. Effect of chlorophyll on lipid oxidation of rapeseed oil. *European Journal of Lipid Science and Technology* 121(4): 1800078. <https://doi.org/10.1002/EJLT.201800078>.
- Machado M, Rodriguez-Alcalá LM, Gomes AM, Pintado M. 2022. Vegetable oils oxidation: mechanisms, consequences and protective strategies. *Food Reviews International* 1–18. <https://doi.org/10.1080/87559129.2022.2026378>.
- Maqsood S, Benjakul S, Abushelaibi A, Alam A. 2014. Phenolic compounds and plant phenolic extracts as natural antioxidants in prevention of lipid oxidation in seafood: a detailed review. *Comprehensive Reviews in Food Science and Food Safety* 13(6): 1125–1140. <https://doi.org/10.1111/1541-4337.12106>.
- Mazaheri Y, Torbati M, Azadmard-Damirchi S, Savage GP. 2019a. Effect of roasting and microwave pre-treatments of *Nigella sativa* L. seeds on lipase activity and the quality of the oil. *Food Chemistry* 274: 480–486. <https://doi.org/10.1016/J.FOODCHEM.2018.09.001>.
- Mazaheri Y, Torbati M, Azadmard-Damirchi S, Savage GP. 2019b. Oil extraction from blends of sunflower and black cumin seeds by cold press and evaluation of its physicochemical properties. *Journal of Food Processing and Preservation* 43(10): e14154. <https://doi.org/10.1111/JFPP.14154>.
- Musial C, Kuban-Jankowska A, Gorska-Ponikowska M. 2020. Beneficial properties of green tea catechins. *International Journal of Molecular Sciences* 21(5): 1744. <https://doi.org/10.3390/IJMS21051744>.
- Naebi M, Torbati M, Azadmard-Damirchi S, Siabi S, Savage GP. 2022. Changes in physicochemical properties of cold press extracted oil from Balangu (*Lallemantia peltata*) seeds during storage. *Journal of Food Composition and Analysis* 107: 104358. <https://doi.org/10.1016/J.JFCA.2021.104358>.
- Nain CW, Berdal G, Thao PTP, et al. 2021. Green tea extract enhances the oxidative stability of DHA-rich oil. *Antioxidants* 10(6): 982. <https://doi.org/10.3390/antiox10060982>.
- Osanloo M, Jamali N, Nematollahi A. 2021. Improving the oxidative stability of virgin olive oil using microformulated vitamin-C. *Food Science & Nutrition* 9(7): 3712–3721. <https://doi.org/10.1002/FSN3.2332>.
- Palit S, Ghosh BC, Dutta Gupta S, Swain DK. 2008. Studies on tea quality grown through conventional and organic management practices: its impact on antioxidant and antidiarrhoeal activity. *Transactions of the ASABE* 51(6): 2227–2238. <https://doi.org/10.13031/2013.25376>.
- Paul A, Radhakrishnan M. 2020. Pomegranate seed oil in food industry: extraction, characterization, and applications. *Trends in Food Science and Technology* 105: 273–283. <https://doi.org/10.1016/j.tifs.2020.09.014>.
- Piravi-Vanak Z, Azadmard-Damirchi S, Kahrizi D, et al. 2022. Physicochemical properties of oil extracted from camelina (*Camelina sativa*) seeds as a new source of vegetable oil in different regions of Iran. *Journal of Molecular Liquids* 345: 117043. <https://doi.org/10.1016/j.molliq.2021.117043>.
- Rezvankhah A, Emam-Djomeh Z, Safari M, Askari G, Salami M. 2018. Investigation on the extraction yield, quality, and thermal properties of hempseed oil during ultrasound-assisted extraction: a comparative study. *Journal of Food Processing and Preservation* 42(10). <https://doi.org/10.1111/jfpp.13766>.
- Rezvankhah A, Emam-Djomeh Z, Safari M, Askari G, Salami M. 2019. Microwave-assisted extraction of hempseed oil: studying and comparing of fatty acid composition, antioxidant activity, physicochemical and thermal properties with Soxhlet extraction. *Journal of Food Science and Technology* 56(9): 4198–4210. <https://doi.org/10.1007/s13197-019-03890-8>.
- Rezvankhah A, Emam-Djomeh Z, Askari G. 2020. Encapsulation and delivery of bioactive compounds using spray and freeze-drying techniques: a review. *Drying Technology* 38(1–2): 235–258. <https://doi.org/10.1080/07373937.2019.1653906>.
- Rezvankhah A, Emam-Djomeh Z, Safari M, Salami M, Askari G. 2022. Investigating the effects of maltodextrin, gum Arabic, and whey protein concentrate on the microencapsulation efficiency and oxidation stability of hemp seed oil. *Journal of Food Processing and Preservation*: e16554. <https://doi.org/10.1111/jfpp.16554>.
- Roshanak S, Rahimmalek M, Goli SAH. 2016. Evaluation of seven different drying treatments in respect to total flavonoid, phenolic, vitamin C content, chlorophyll, antioxidant activity and color of green tea (*Camellia sinensis* or *C. assamica*) leaves. *Journal of Food Science and Technology* 53(1): 721–729. <https://doi.org/10.1007/s13197-015-2030-x>.
- Rostami M, Farzaneh V, Boujmehrani A, Mohammadi M, Bakhshabadi H. 2014. Optimizing the extraction process of sesame seed's oil using response surface method on the industrial scale. *Industrial Crops and Products* 58: 160–165. <https://doi.org/10.1016/J.INDCROP.2014.04.015>.
- Sahin S, Sayim E, Bilgin M. 2017. Effect of olive leaf extract rich in oleuropein on the quality of virgin olive oil. *Journal of Food Science and Technology* 54(6): 1721–1728. <https://doi.org/10.1007/S13197-017-2607-7>.
- Samadi S, Raouf Fard F. 2020. Phytochemical properties, antioxidant activity and mineral content (Fe, Zn and Cu) in Iranian produced black tea, green tea and roselle calyces. *Biocatalysis and Agricultural Biotechnology* 23: 101472. <https://doi.org/10.1016/J.BCAB.2019.101472>.
- Tian Y, Xu Z, Zheng B, Martin Lo Y. 2013. Optimization of ultrasonic-assisted extraction of pomegranate (*Punica granatum* L.) seed oil. *Ultrasonics Sonochemistry* 20(1): 202–208. <https://doi.org/10.1016/j.ultsonch.2012.07.010>.
- Unno K, Furushima D, Hamamoto S, et al. 2018. Stress-reducing function of matcha green tea in animal experiments and clinical trials. *Nutrients* 10(10). <https://doi.org/10.3390/nu10101468>.

- Wang Y, Zhou S, Jiang Y, *et al.* 2021. Effects of tea polyphenol palmitate on the crystallization of rice bran oil. *Journal of Food Engineering* 292: 110279. <https://doi.org/10.1016/J.JFOODENG.2020.110279>.
- Womani HM, Djikeng FT, Anjaneyulu B, Karuna MSL, Prasad RBN, Linder M. 2016. Oxidative stabilization of RBD palm olein under forced storage conditions by old Cameroonian green tea leaves methanolic extract. *NFS Journal* 3: 33–40. <https://doi.org/10.1016/j.nfs.2016.03.002>.
- Ye Z, Zhuang Y, Chen Y, *et al.* 2020. Aqueous-phase oxidation of three phenolic compounds by hydroxyl radical: insight into secondary organic aerosol formation yields, mechanisms, products and optical properties. *Atmospheric Environment* 223: 117240. <https://doi.org/10.1016/J.ATMOSENV.2019.117240>.
- Yeasmen N, Orsat V. 2021. Green extraction and characterization of leaves phenolic compounds: a comprehensive review. *Crit Rev Food Sci Nutr* 1–39. <https://doi.org/10.1080/10408398.2021.2013771>.

Cite this article as: Dezashibi Z, Azadmard-Damirchi S, Piravi-Vanak Z. 2022. Effect of co-extraction of pomegranate seed oil with green tea leaves on the extraction yield and quality of extracted oil. *OCL* 29: 25.